

Amendments to the Specification

Please replace the paragraph beginning at page 7, lines 1-11, with the following rewritten paragraph:

include imipramine, amitriptyline, nortriptyline, and desipramine. Monoamine oxidase inhibitors (MAOIs) are also used as antidepressants. MAOIs approved for the treatment of depression include phenelzine (~~Nardil~~ NARDIL®), tranylcypromine (PARNATE® ~~Parnate~~), and isocarboxazid (MARPLAN® ~~Marplan~~). Medications that primarily affect the neurotransmitter serotonin, termed selective serotonin reuptake inhibitors (SSRIs), are also used as antidepressants. These include escitalopram HBr (LEXAPRO® ~~Lexapro~~), fluoxetine (PROZAC® ~~Prozac~~), sertraline (ZOLOFT® ~~Zoloft~~), fluvoxamine (LUVOX® ~~Luvox~~), paroxetine (PAXIL® ~~Paxil~~), and citalopram (CELEXA® ~~Celexa~~). Additional medication of use affect both norepinephrine and serotonin, for example venlafaxine (~~Effexor~~ EFFEXOR®) and nefazadone (SERZONE® ~~Serzone~~), or agents such as phenelzine (~~Nardil~~ NARDIL®), tranylcypromine (PARNATE® ~~Parnate~~), mirtazepine (REMERON® ~~Remeron~~), nefazodone (SERZONE® ~~Serzone~~), triazolopyridine (TRAZODONE® ~~trazodone~~), and bupropion (WELLBUTRIN® ~~Wellbutrin~~).

Please replace the paragraphs on page 17, lines 5-8 with the following rewritten paragraph:

Example 1

Patient 1

A 59-year-old Caucasian female with a history of depression for 3 years was treated. Medications included PAXIL® ~~Paxil~~ (20 mg), as prescribed by her psychiatrist. She appeared somewhat depressed at the time of the first office visit. Her glabellar frown lines and forehead lines were treated with 40 i.u. of BOTOX™ (Fig. 4).

Please replace the paragraph at page 17, lines 22-27, with the following rewritten paragraph:

Example 2

Patient 2

A 33-year-old Caucasian female with a history of intermittent anxiety/depression for ten years was treated. Her medications included an SSRI (ZOLOFT® Zoloft, 200 mg per day) for the last four years. Prior to treatment she reported feeling depressed and anxious.

Please replace the paragraph at page 18, lines 1-10 with the following rewritten paragraph:

Example 3

Patient 3

A 50-year-old, recently divorced, Caucasian female was treated. She had a history of depression for two years. Her medications included PAXIL® Paxil-20 mg orally per day and Wellbutrin 225 mg orally per day. She reported feeling depressed.

Her frown/forehead lines were injected with 30 Unit equivalents of BOTOX™ (Fig. 6). Four months later she reported a much-improved mood, and stated that she did not feel depressed. Her frown/forehead lines were re-treated with 30 Unit equivalents of BOTOX™.